

Reconstructing the interaction between the dark matter and holographic dark energy

Shao-Feng Wu^{1,2*}, Peng-Ming Zhang^{3,4}, and Guo-Hong Yang^{1,2}

¹*Department of Physics, College of Science, Shanghai University, Shanghai, 200444, P. R. China*

²*The Shanghai Key Lab of Astrophysics, Shanghai, P. R. China*

³*Center of Theoretical Nuclear Physics, National Laboratory of Heavy Ion Accelerator, Lanzhou 730000, P. R. China and*

⁴*Institute of Modern Physics, Lanzhou, 730000, P. R. China*

We reconstruct the interaction rate between the dark matter and the holographic dark energy with the parameterized equation of states and the future event horizon as the infrared cut-off length. It is shown that the observational constraints from the 192 SNIa and BAO measurement permit the negative interaction in the wide region. Moreover, the usual phenomenological descriptions can not describe the reconstructed interaction well for many cases. The other possible interaction is also discussed.

PACS numbers: 95.36.+x, 98.80.-k, 98.80.Cq

I. INTRODUCTION

In the modern cosmology, a ‘dark energy’ (DE) with negative pressure is suggested to be responsible for the current acceleration of the universe. The simplest candidate of DE is the cosmological constant, which does nicely well at the pragmatic observational level, but entails the serious theoretical difficulty: the cosmological constant problem and the coincidence problem. Explanations of DE have been sought within a wide range of physical phenomena, including some exotic fields, modified gravity theories, and so on -see [1] and references therein. Among the most recent generic proposals, the model inspired by the holographic idea [2], that the quantum zero-point energy of a system cannot exceed the mass of a black hole with same size, has been put forward to explain the DE [3, 4, 5]. This DE density can be determined in terms of the horizon radius of the universe, corresponding to relate the UV cutoff of a system to its IR cutoff in the quantum field theory. There are usually three choices for the horizon radius supposed to provide the IR cutoff, with different degrees of success, namely the Hubble horizon, the particle horizon, and the future event horizon. The event horizon may be better, since in this case the DE can drive the present accelerated expansion and the coincidence problem can be resolved by assuming an appropriate number of e-folding of inflation [5, 6, 7].

Most discussions on DE models rely on the fact that both dark matter (DM) and DE only couple gravitationally. However, given their unknown nature and the symmetry that would impose a vanishing interaction is still to be discovered, an entirely independent behavior between dark sectors is very special. Moreover, since DE must be accreted by massive compact objects like black holes and neutron stars, in a cosmological context the energy transfer from DE to DM may be small but must be non-vanishing. The interaction hypothesis was first introduced by Wetterich [8] to discuss the cosmological constant problem in the light of dilatation symmetry and its anomaly. Then cosmological consequences of a scalar field coupled to the matter were studied in [9]. It was found that the coupling quintessence models may give the scaling attractors providing an accelerated expansion at the present time and alleviate the coincidence problem [10]. The interaction also appears in the context of modified gravity models [11]. More possibility that DE and DM can interact has been studied in [12, 13, 14, 15, 16, 17]. Confronted to cosmological data, it was found that an appropriate interaction can influence the perturbation dynamics, and the lowest multipoles of the CMB spectrum [18, 19], and could be inferred from the expansion history of the Universe, as manifested in the supernova data together with CMB and large-scale structure [20, 21, 22]. In addition, it was suggested that the dynamical equilibrium of collapsed structures would be affected by the coupling of DE to DM [23, 24]. The interaction was first connected to holography by Horvat [25] who argued that scaling of the cosmological constant stemming from the zero-point energy in quantum field theory possibly implies a non-vanishing coupling of the cosmological constant with DM. In the holographic DE model with the Hubble horizon as the IR cutoff, the interaction can be available to derive the present accelerated expansion and alleviate the coincidence problem [26]. In the interacting model with the event horizon as the IR cutoff, it was shown that the equation of state (EoS) of DE can accommodate the dynamically evolving behavior of crossing the phantom divide [27], which suggested by recent most observational probes [28].

Although the interaction is important in studying the physics of DE, it will not be possible to derive the precise form of the interaction from first principles unless the nature of both dark sectors was known. Usually, the coupling is determined from phenomenological requirements [10, 26]. In view of the continuous equations of DE density ρ_d and

* Corresponding author. Email: sfwu@shu.edu.cn; Phone: +86-021-66136202.

DM density ρ_m , the coupling Q must be a function of densities multiplied by a quantity with units of inverse of time, which has an obvious choice as Hubble time H^{-1} . Thus, one may write the coupling as

$$Q = Q(H\rho_m, H\rho_d), \quad (1)$$

which leads $Q \simeq \lambda_m H\rho_m + \lambda_d H\rho_d$ from the first order terms in the power law expansion. Assuming that the ratio $r = \rho_m/\rho_d$ might be piecewise constant, the linear parameters are usually set to $\lambda_m \simeq \lambda_d$ and even $\lambda_m \simeq 0$ or $\lambda_d \simeq 0$ for simplicity. Considering the couplings are terms in the Lagrangian which mix both DE and DM, one may further suppose that they could be parameterized by some product of the densities of DE and DM, such as the simplest $Q \simeq \lambda\rho_m\rho_d$ [13]. Besides these phenomenological descriptions, various proposals at the fundamental level have been tried to account for the coupling, including the dependence of the matter field on the scalar field [29] or expressing the cosmological constant as a function of the trace of the energy-momentum tensor [30]. Recently, an interesting thermodynamical description of interaction between holographic DE and DM has been proposed in [31], where it was assumed that in the absence of the coupling the DE and DM remain in separate thermal equilibrium, then a small interaction can be viewed as a stable thermal fluctuation that brings a logarithmic correction to the equilibrium entropy of DE and DM. Other specific coupling which was assumed from the outset can be found in [12, 32, 33].

The main aim of this work is to reconstruct the coupling using the recent DE probes (the Baryon Acoustic Oscillation (BAO) measurement at $z = 0.35$ from the Sloan Digital Sky Survey [34] and the re-compiled 192 Type Ia Supernovae (SnIa) samples [35], consisting 60 points from ESSENCE (“Equation of State: Supernovae trace Cosmic Expansion”) supernova survey [36], 57 points from Supernovae Legacy Survey [37], 45 points nearby Supernovae [38], and 30 points detected by the Hubble Space Telescope [39]). We will focus on the holographic DE model and choose the future event horizon as the IR cutoff.

The model with the Hubble horizon has been studied recently in [40] and it was found that the reconstructed interaction is always positive in 1σ region. This seems to corroborate the recent argument that the negative interaction violates the second law of thermodynamics which inquires the energy transfer from DE to DM rather than otherwise [41]. However, it should be noticed that there are some problems in the thermodynamics of DE, such as the negative entropy [42], and the generalized thermodynamical second law indeed breaks down in the universe with phantom-dominated DE [43, 44]. These results suggest that one should consider the thermodynamical properties of DE with wide possibilities. Hence, it is interesting to see whether the positive interaction is a robust result for other models, such as the present model with the IR cutoff as the future event horizon.

Considering the time varying DE gives a better fit than a cosmological constant and in particular most of the observational probes indeed mildly favor dynamical DE crossing the phantom divide at $z \sim 0.2$ [28], we will employ two commonly used parameterizations [21, 45, 46, 47], namely

$$w_A = w_0 + w_1(1 - a) = w_0 + w_1 \frac{z}{1 + z}, \quad (2)$$

which has been used in [40], and

$$w_B = w_0 + w_1(1 - a)a = w_0 + w_1 \frac{z}{(1 + z)^2}. \quad (3)$$

It should be noticed that the different parameterizations are beneficial to control some amount of parameterization dependence. After reconstructing the interaction, we will further compare it with the usual phenomenological models and the recent thermodynamical description.

II. RECONSTRUCTION

Let us begin with the Friedmann equations

$$H^2 = \frac{1}{3}(\rho_m + \rho_d), \quad (4)$$

$$\dot{H} = -\frac{1}{2}(\rho_m + \rho_d + p_m + p_d), \quad (5)$$

where we have normalized $8\pi G = 1$ for conventions. The total energy density $\rho = \rho_m + \rho_d$ satisfies a conservation law. However, since we consider the interaction between DE and DM, ρ_m and ρ_d do not satisfy independent conservation laws, they instead satisfy two continuous equations

$$\dot{\rho}_m + 3H\rho_m = Q, \quad (6)$$

$$\dot{\rho}_d + 3H(1 + w_d)\rho_d = -Q, \quad (7)$$

where w_d is the EoS of DE, and Q denotes the interaction term. Without loss of generality, we will write the interaction as $Q = \rho_d \Gamma$, where Γ is an unknown function.

Using the the ratio of energy densities $r = \rho_m/\rho_d$, we have

$$\frac{\dot{H}}{H^2} = -\frac{3}{2} \frac{1 + w_d + r}{1 + r}. \quad (8)$$

from Eq. (5). From Eq. (6), we have

$$\dot{r} = -r \frac{\dot{\rho}_d}{\rho_d} - 3Hr + \Gamma,$$

which can be recast as

$$\dot{r} = (1 + r)\Gamma + 3Hrw_d \quad (9)$$

from Eq. (7). Eliminating the Γ in above two equations, we obtain

$$\frac{-r\dot{r} - 3Hrw_d}{1 + r} = r \frac{\dot{\rho}_d}{\rho_d} + 3Hr. \quad (10)$$

Until now, we have not specified the concrete DE density. We will focus on the holographic DE model. Followed [5] by choosing the future event horizon

$$R_E = a \int_a^\infty \frac{dx}{Hx^2} \quad (11)$$

as the IR cutoff, the holographic DE density is $\rho_d = 3c^2 R_E^{-2}$, where c^2 is a constant and the Planck mass has been taken as unit. The most possible theoretical value of c is one [5, 6], indicating that the total energy from DE must be determined by the Schwarzschild relation. Taking the derivative with respect to t , the evolution of the horizon can be determined by

$$\dot{R}_E = HR_E - 1.$$

Defining $\Omega_d = \rho_d/3H^2$, we have $\Omega_d = 1/(1 + r)$ and $R_E = c\sqrt{1 + r}/H$. Thus, Eq. (10) can be recast as

$$\frac{-r\dot{r} - 3Hrw_d}{1 + r} = r \frac{-2H(c\sqrt{1 + r} - 1)}{c\sqrt{1 + r}} + 3Hr. \quad (12)$$

Obviously, r is not a constant in general. This is different with the case in [40] where r is a constant since the IR cutoff was chosen as being the Hubble scale. Replacing the time t as the redshift $z = (1 - a)/a$, we can rewrite Eq. (12) as

$$\frac{-(1 + z)rr' + 3rw_d}{1 + r} = r \frac{2(c\sqrt{1 + r} - 1)}{c\sqrt{1 + r}} - 3r, \quad (13)$$

where the prime denotes the derivative with respect to z . Similarly, Eq. (8) reads

$$\frac{H'}{H} = \frac{3}{2(1 + z)} \frac{1 + w_d + r}{1 + r}. \quad (14)$$

It is interesting to find that Eqs. (13) and (14) determine the evolvement of r and H , if we know the EoS w_d . In the normal interacting DE model, one assumes the explicit interaction form to give out w_d . For our aim, we will use the two commonly used parameterizations of w_d to determine the dynamics of our model, and reconstruct the interaction rate $\Gamma/3H$

$$\frac{\Gamma}{3H} = -\frac{(1 + z)r' + 3rw_d}{3(1 + r)}$$




FIG. 1: The deceleration parameter q with the best-fit values of equation of states w_d . The red, green, and blue lines denote the models without interaction, with interaction and $w_d = w_A$, and with interaction and $w_d = w_B$, respectively. For each model, there are five lines from top to down at early taking different c from 0.6 to 1.4.

from the recent observational datasets.

We will use the re-compiled 192 SnIa samples ($0 < z < 1.8$) combined with the recent BAO measurement from SDSS to reconstruct the interaction rate. There are other DE observational probes, including the three-year WMAP CMB shift parameter, the X-ray gas mass fraction in clusters, the linear growth rate of perturbations at $z = 0.15$ as obtained from the 2dF galaxy redshift survey, and the look back age data. However, since the parameterizations of w_d Eqs. (2) and (3) are motivated to accommodate the dynamically evolving behavior of crossing the phantom divide at recent epoch, and our model has not included the radiation and the baryonic matter which may be important in the early, we will not use the WMAP CMB shift parameter which focuses on the high redshift region. Besides, for simplicity, we do not adopt other probes of DE which have large relative errors compared with SnIa, CMB, and BAO probes [48].

As usually, we will fix DM density parameter as $\Omega_{0m} = 0.3$ or 0.25 to include the best-fit value of $\Omega_{0m} = 0.27$ from 5-year WMAP data. In general, they are sufficiently representative. Moreover, it is convenient to compare our reconstructed interaction to the interaction reconstructed in [40] where these two DM density parameters are used. We will consider five indicative different values of the constant c near one. The best-fit values with 1σ error bars for the parameters w_0 and w_1 are given in TABLE I, II. One can find that the phantom divide crossing in recent

	c	0.6	0.8	1.0	1.2	1.4
w_A	w_0	-1.19 ± 0.18	-1.16 ± 0.16	-1.14 ± 0.16	-1.13 ± 0.15	-1.12 ± 0.14
	w_1	0.60 ± 1.26	1.09 ± 1.07	1.35 ± 0.97	1.50 ± 0.90	1.60 ± 0.85
w_B	w_0	-1.21 ± 0.24	-1.21 ± 0.22	-1.21 ± 0.20	-1.21 ± 0.19	-1.20 ± 0.19
	w_1	0.99 ± 2.11	1.83 ± 1.83	2.29 ± 1.66	2.58 ± 1.55	2.78 ± 1.47

TABLE I: The best fit values with 1σ error bars for the parameters of w_d with DM density parameter $\Omega_{0m} = 0.25$.

	c	0.6	0.8	1.0	1.2	1.4
w_A	w_0	-1.03 ± 0.20	-1.03 ± 0.18	-1.03 ± 0.16	-1.03 ± 0.16	-1.02 ± 0.15
	w_1	0.48 ± 1.28	0.48 ± 1.09	0.48 ± 0.98	0.70 ± 0.92	0.85 ± 0.87
w_B	w_0	-0.99 ± 0.26	-1.02 ± 0.23	-1.04 ± 0.22	-1.05 ± 0.21	-1.05 ± 0.20
	w_1	-1.14 ± 2.20	0.03 ± 1.90	0.68 ± 1.73	1.09 ± 1.62	1.37 ± 1.54

TABLE II: The best fit values with 1σ error bars for the parameters of w_d with DM density parameter $\Omega_{0m} = 0.3$.

epoch is always permitted in 1σ region. The deceleration parameter with these best-fit equation of states is plotted in FIG. 1, where the acceleration (super-acceleration, at most cases) in recent epoch is achieved. It is interesting to see that current deceleration parameter in interacting model is almost not affected by the DE parameter c , according to the almost same parameters w_0 in TABLE I, II. We also show the evolution of ρ_m compared its non-interacting version ρ_m^0 in FIG. 2. The ratios change strongly in recent epoch which implies the interaction plays important role




FIG. 2: The ratios ρ_m/ρ_m^0 for different Ω_{0m} , w_d , and c . In each panel, c increases from top to bottom.




FIG. 3: The ratios ρ_d/ρ_d^0 for different Ω_{0m} , w_d , and c . In each panel, c increases from top to bottom.

in that time. The corresponding plots for the dark energy component are given in FIG. 3. Since the DE density is always positive, this suggests that the interaction can be negative but can not be too negative when $w_d > -1$, which is possible at least in the past.

With these best-fit values, we reconstruct the interaction rates, see FIG. 4 and FIG. 5. This is one of the main result of this paper. It should be stressed that the negative interaction is permitted in the wide region of FIG. 4 and FIG. 5. This is contrast to the result in [40] where the reconstructed interaction is always positive, and the argument that the second law of thermodynamics which imposes energy transfer from DE to DM [41]. If the second law is true, the permitted region of the interaction rates must be reduced, as done in [49]. However, for generality and considering the thermodynamical properties of DE is not clear, we will not restrict the interaction to be positive.

Moreover, the effective EoS $w_{eff} = w_d + \frac{\Gamma}{3H}$ can be obtained since we have known w_d and reconstructed the $\Gamma/3H$. From TABLE I, II, and FIG. 4, 5, one can know that the effective EoS may be bigger or smaller than -1 for different parameters Ω_{0m} and c . If one requires an effective phantom-like DE, the presence of a coupling makes the parameters have more possible values than the case of absence of a coupling, where the EoS is determined by $w_d = -\frac{1}{3} - \frac{2}{3c}\sqrt{\Omega_d}$ which imposes $c < \sqrt{\Omega_d}$ for $w_d < -1$. Moreover, it is possible to have effective EoS w_{eff} smaller than -1 but w_d bigger than -1 when the interaction is negative. This suggests that the negative interaction is interesting because DE with $w_d > -1$ is easily accepted (The scalar field with $w_d < -1$ will break the zero energy condition.) and the effective EoS of DE $w_{eff} < -1$ can fit the cosmological data better.

III. COMPARISON

In the following, we will compare the reconstructed interaction rate with other descriptions. Let us begin with the usual phenomenological descriptions. Usually there are four different choices of Q_i ($i = 1, 2, 3, 4$), which can be expressed as $3\lambda H\rho_d$, $3\lambda H\rho_m$, $3\lambda H(\rho_m + \rho_d)$, and $\lambda\rho_m\rho_d$, respectively. To compare them with the reconstructed interaction rate $\Gamma/3H$, we define four interaction rates $R_i = Q_i/(3H\rho_d)$, which can be determined by r and H (which have been solved from Eqs. (13) and (14)), namely $R_1 = \lambda$, $R_2 = \lambda r$, $R_3 = \lambda(1+r)$, and $R_4 = \lambda Hr/(1+r)$. Observing FIG. 4, the indicative curves of the usual interaction rates R_i ($i = 1, 2, 3, 4$) show that in many cases they are not favored in 1σ region. For example, see the case of $c = 1.2$, where the interaction rates R_2 (green line) and R_3 (blue line) are permitted in 1σ region, but the other interaction rates R_1 (red line) and R_4 (yellow line) are not permitted whether λ increases or decreases. For the case $c = 0.6$, the reconstructed interaction rate changes at recent epoch




FIG. 4: Interaction rates for $\Omega_{0m} = 0.3$ with respect to redshift z . Above and below panels denote $w_d = w_A$ and w_B , respectively. The black curves and the grey region between them indicate the best fit curve of the reconstructed interaction rate $\Gamma/3H$ and the region in 1σ confidence level. The black dashed lines indicate $\Gamma/3H = 0$. The red, green, blue, and yellow curves indicate the R_i from $i = 1$ to $i = 4$ with four constants λ . For example, we give $\lambda = 0.13, 0.11, 0.07, 0.2$ in turn for $w_d = w_A$ and $c = 1$. The blue dashed lines indicative the interaction rate R_5 . The red dashed lines in the case with $c = 0.6$ indicative the interaction rate R_6 with two nonzero constants. For the above panel, they are $\lambda_1 = 0.17, \lambda_2 = -0.16$; For the below panel, they are $\lambda_1 = 0.11, \lambda_2 = -0.15$.




FIG. 5: Interaction rates for $\Omega_{0m} = 0.25$. The blue dashed lines indicative the interaction rate R_5 . The red dashed lines indicative the interaction rate R_6 with some constants λ_i ($i = 1, \dots, 5$).

from positive to negative. This tendency is more general in FIG. 5 with small DM parameter $\Omega_{0m} = 0.25$. Thus, one may suspect that the four phenomenological descriptions of the interaction which have the definitive sign may be not suitable.

We will also consider the recent thermodynamical description of the interaction. Defining $Q_5 = 3Hb^2(\rho_m + \rho_d)$, the thermodynamical interaction rate b^2 can be determined by following equations (see [31] in detail):

$$b^2 = \frac{2\Omega_d^{3/2}}{3c} \left[-1 + \frac{H^2\sqrt{\Omega_d}}{(H^0)^2\sqrt{\Omega_d^0}} \frac{\sqrt{\Omega_d^0/c} - 1}{\sqrt{\Omega_d/c} - 1} \right] + \frac{1}{12\pi c^2} \frac{H^2}{c/\sqrt{\Omega_d}(c/\sqrt{\Omega_d} - 1)} \frac{\sqrt{\Omega_d}}{\sqrt{\Omega_d^0}} (-\Omega_d^0)'(1+z),$$

$$-(1+z)\frac{\Omega_d'}{\Omega_d} + (\Omega_d - 1) + \frac{2\sqrt{\Omega_d}}{c}(\Omega_d - 1) = -3b^2,$$

$$\frac{-(1+z)H'}{H} = \frac{\sqrt{\Omega_d}}{c} - 1 + \frac{(1+z)\Omega_d'}{2\Omega_d},$$

where the zero superscript of Ω_d^0 and H^0 indicates absence of interaction. We can relate b^2 to a new interaction rate

$$R_5 = \frac{Q_5}{3H\rho_d} = \frac{b^2(\rho_m + \rho_d)}{\rho_d} = \frac{b^2}{\Omega_d}.$$

In FIG. 4 and FIG. 5 (blue dashed lines), one can find that R_5 is almost not favored in 1σ region (Note we have not shown the cases with $c < 1$ where b^2 has not the real solution and the parameterization of w_d is not needed to determine R_5 since the EoS in this model is determined by the b^2).

It is interesting to ask whether the reconstructed interaction can be described well by a more general interaction form. A natural candidate is to expand the phenomenological description (1) up to the second order, namely

$$Q_6 = 3\lambda_1 H\rho_m + 3\lambda_2 H\rho_d + \lambda_3 H^2\rho_m\rho_d + \lambda_4 H^2\rho_m^2 + \lambda_5 H^2\rho_d^2,$$

with the interaction rate

$$R_6 = \frac{Q_6}{3H\rho_d} = \lambda_1 r + \lambda_2 + \lambda_3 \frac{H^3 r}{1+r} + \lambda_4 \frac{H^3 r^2}{1+r} + \lambda_5 \frac{H^3}{1+r}.$$

We find that it indeed works well, see the red dashed line in FIG. 4 and FIG. 5. However, this parameterization only has theoretical interest since it contains too many parameters. Moreover, numerical calculations prove that this parameterization can not be reduced to include only the two first order terms, which has been studied recently in [49], if we need it being permitted in 1σ region for all cases.

IV. SUMMARY

We have reconstructed the interaction term between the holographic DE and DM, using the re-compiled 192 SnIa samples combined with the recent BAO measurement. The DE parameter c is assumed near one. The two common used parameterizations of w_d are considered in 1σ region. It is found that the present accelerated expansion of universe is achieved and the phantom behavior of DE is permitted. We illustrate that the negative interaction is permitted in the wide region. Hence, contrast to the DE model studied in [40], where the reconstructed interaction is always positive in 1σ region, we can not obtain the favor from the DM observation for the recent argued thermodynamical second law which imposes the energy transfer from DE to DM. This suggest us to keep wide possibilities of the thermodynamical properties of DE.

We show that the four usual phenomenological descriptions can not describe the reconstructed interaction well for many cases. Specially for the case with small DM parameter $\Omega_{0m} = 0.25$, the interaction rate has a trend to change sign at recent epoch. This is at variance with the usual phenomenological interacting terms which have the definitive sign. We further illustrate that the recent thermodynamical description of the interaction is not favored. Our work stimulates one to seek a more suitable interaction.

It would be interesting to confront our model to more observations, such as CMB angular power and large scale structure, which will further constrain the interaction rate and make clear if the negative interaction and phantom DE are still to be permitted. Another interesting work is to consider the perturbation evolution in our model. Recently, it has been found that some types of interaction will lead to instability under the curvature perturbation [50, 51]. It may provide some restrictions on the form of interaction. We will work on these directions in the future.

Acknowledgments

This work was supported by NSFC under Grant Nos. 10847102 and 10604024, the Shanghai Research Foundation No. 07dz22020, the CAS Knowledge Innovation Project Nos. KJcx.syw.N2, the Shanghai Education Development Foundation, the Innovation Foundation of Shanghai University, and Shanghai Leading Academic Discipline Project S30105.

-
- [1] S. Padmanabhan, Phys. Rep. **380**, 235 (2003); P. J. E. Peebles, and B. Ratra, Rev. Mod. Phys. **75**, 559 (2003); L. Perivolaropoulos, astro-ph/0601014.
 - [2] G. 't Hooft, gr-qc/9310026; L. Susskind, J. Math. Phys. **36**, 6377 (1995).
 - [3] A. Cohen, D. Kaplan, A. Nelson, Phys. Rev. Lett. **82** (1999) 4971.
 - [4] S. D. H. Hsu, Phys. Lett. **B 594**, 13 (2004).
 - [5] M. Li, Phys. Lett. **B 603**, 1 (2004); Q. G. Huang and M. Li, JCAP **0408**, 013 (2004).
 - [6] M. Li, C. Lin, Y. Wang, JCAP **05**, 023 (2008).
 - [7] J. Lee, H.C. Kim, J.W. Lee, Phys. Lett. **B 661**, 67 (2008).
 - [8] C. Wetterich, Nucl. Phys. **B 302**, 668 (1988).
 - [9] J. Ellis, S. Kalara, K. A. Olive, and C. Wetterich, Phys. Lett. **B 228**, 264 (1989); T. Damour and K. Nordtvedt, Phys. Rev. **D 48**, 3436 (1993).
 - [10] L. Amendola, Phys. Rev. **D 62**, 043511 (2000); W. Zimdahl, D. Pavón, and L. P. Chimento, Phys. Lett. **B 521**, 133 (2001); D. Tocchini-Valentini, and L. Amendola, Phys. Rev. **D 65**, 063508 (2002); L. P. Chimento, A. S. Jakubi, D. Pavón, and W. Zimdahl, Phys. Rev. **D 67**, 083513 (2003); S. del Campo, R. Herrera and D. Pavón, Phys. Rev. **D 70**, 043540 (2004); G. Olivares, F. Atrio-Barandela, D. Pavón, Phys. Rev. **D 71**, 063523 (2005); L. P. Chimento and D. Pavón, Phys. Rev. **D 73**, 063511 (2006); D. F. Mota and D. J. Shaw, Phys. Rev. **D 75**, 063501 (2007); G. Olivares, F. Atrio-Barandela, D. Pavon, Phys. Rev. **D 77**, 063513 (2008).
 - [11] L. Amendola, Phys. Rev. **D 60**, 043501 (1999); L. Amendola, D. Polarski, and S. Tsujikawa, Phys. Rev. Lett. **98**, 131302 (2007); N. Banerjee, D. Pavón, Phys. Lett. **B 647**, 477 (2007); S. Das, N. Banerjee, Phys. Rev. **D 78**, 043512, (2008).
 - [12] G. W. Anderson, S. M. Carroll, astro-ph/9711288.
 - [13] G. Mangano, G. Miele and V. Pettorino, Mod. Phys. Lett. **A 18**, 831 (2003).
 - [14] G. R. Farrar and P. J. E. Peebles, Astrophys. J. **604**, 1 (2004).
 - [15] E. Elizalde, S. Nojiri, S. D. Odintsov, and P. Wang, Phys. Rev. **D 71**, 103504 (2005).
 - [16] M. R. Setare, Phys. Lett. **B 642** (2006) 1; Phys. Lett. **B 642**, 421 (2006).
 - [17] S. B. Chen, B. Wang, J. J. Jing, Phys. Rev. **D 78**, 123503, (2008).
 - [18] W. Zimdahl, Int. J. Mod. Phys. **D 14**, 2319 (2005).
 - [19] B. Wang, J. Zang, Ch.-Y. Lin, E. Abdalla, S. Micheletti, Nucl. Phys. **B 778**, 69 (2007).
 - [20] Z. K. Guo, N. Ohta, and S. Tsujikawa, Phys. Rev. **D 76**, 023508 (2007).
 - [21] C. Feng, B. Wang, Y. Gong, R.-K. Su, JCAP **09**, 005 (2007); J. H. He, B. Wang, JCAP **06**, 010 (2008).
 - [22] L. Amendola, G. Campos, R. Rosenfeld, Phys. Rev. **D 75**, 083506 (2007).
 - [23] M. Kesden and M. Kamionkowski, Phys. Rev. Lett. **97**, 131303 (2006); M. Kesden and M. Kamionkowski, Phys. Rev. **D 74**, 083007 (2006).
 - [24] O. Bertolami, F. Gil Pedro and M. Le Delliou, Phys. Lett. **B 654**, 165 (2007).
 - [25] R. Horvat, Phys. Rev. **D 70**, 087301 (2004).
 - [26] D. Pavón and W. Zimdahl, Phys. Lett. **B 628**, 206 (2005); W. Zimdahl and D. Pavón, Class. Quantum Grav. **24**, 5461 (2007).
 - [27] B. Wang, Y. Gong, E. Abdalla, Phys. Lett. **B 624**, 141 (2005); B. Wang, Ch.-Y. Lin, E. Abdalla, Phys. Lett. **B 637**, 357 (2006).
 - [28] Y. Wang and M. Tegmark, Phys. Rev. Lett. **92**, 241302 (2004); U. Alam, V. Sahni and A. Starobinsky, JCAP **0406**, 008 (2004); P. S. Corasaniti et al., Phys. Rev. **D 70**, 083006 (2004); S. Hannestad and E. Mortsell, JCAP **0409**, 001 (2004); A. Upadhye, M. Ishak and P. J. Steinhardt, Phys. Rev. **D 72**, 063501 (2005).
 - [29] F. Piazza, S. Tsujikawa, JCAP **07**, 004 (2004).
 - [30] N. J. Poplawski, gr-qc/0608031.
 - [31] B. Wang, C.-Y. Lin, D. Pavón, E. Abdalla, Phys. Lett. **B 662**, 1 (2008).
 - [32] L. Amendola, D. Tocchini-Valentini, Phys. Rev. **D 64**, 043509 (2001).
 - [33] S. Das, P.S. Corasaniti, J. Khoury, Phys. Rev. **D 73**, 083509 (2006).
 - [34] D. J. Eisenstein, *et al.*, Astrophys. J. **633**, 560 (2005).
 - [35] T. M. Davis *et al.*, Astrophys. J. **666**, 716 (2007).
 - [36] W. M. Wood-Vasey *et al.*, Astrophys. J. **666**, 694 (2007); G. Miknaitis *et al.*, astro-ph/0701043.
 - [37] P. Astier *et al.*, Astron. Astrophys. **447**, 31 (2006).
 - [38] A. G. Riess *et al.*, Astrophys. J. **607**, 665 (2004).
 - [39] A. G. Riess *et al.*, astro-ph/0611572.
 - [40] A. A. Sen, and D. Pavón, Phys. Lett. **B 664**, 7 (2008).

- [41] D. Pavón and B. Wang, arXiv:0712.0565.
- [42] I. Brevik, S. Nojiri, S. D. Odintsov, L. Vanzo, Phys. Rev. **D 70**, 043520 (2004); S. Nojiri and S. D. Odintsov, Phys. Rev. **D 72**, 023003 (2005).
- [43] G. Izquierdo, D. Pavón, Phys. Lett. **B 639**, 1 (2006).
- [44] H. M. Sadjadi, Phys. Lett. **B 645**, 108 (2007).
- [45] H. K. Jassal, J. S. Bagla, and T. Padmanabhan, Mon. Not. Roy. Astron. Soc. **356**, L11 (2005).
- [46] M. Chevallier, D. Polarski, Int. J. Mod. Phys. **D 10**, 213 (2001).
- [47] E. V. Linder, Phys. Rev. Lett. **90**, 091301 (2003).
- [48] S. Nesseris and L. Perivolaropoulou, JCAP **0701**, 018 (2007).
- [49] K. Karwan, JCAP **0805**, 011 (2008).
- [50] J. Valiviita, E. Majerotto, R. Maartens, JCAP **0807**, 020, (2008).
- [51] J. H. He, B. Wang, E. Abdalla, arXiv:0807.3471 [gr-qc].